

(place patient label here)

Patient Name: _____



PROVIDER ORDERS

Order Set Directions:

- > (✓)- Check orders to activate; Orders with pre-checked box will be followed unless lined out.
- > Initial each place in the pre-printed order set where changes such as additions, deletions or line outs have been made
- > Initial each page and Sign/Date/Time last page

Diagnosis: _____

Allergies with reaction type: _____

Pain Management Medications

Version 1 Approved 1/16/14

- For Moderate and Severe Pain: May select one oral and/or one injectable medication (not all),

Analgesics

Oral

- acetaminophen (TYLENOL)
 - 650 milligram orally every 4 hours as needed for mild-to-moderate pain
- ibuprofen (MOTRIN)
 - 400 milligram orally every 6 hours as needed for mild-to-moderate pain
- HYDROcodone-acetaminophen 5 mg-325 mg tab (NORCO)
 - 1-2 tablet orally every 4 hours as needed for moderate-to-severe pain
- oxyCODONE-acetaminophen 5 mg-325 mg tab (PERCOCET)
 - 1-2 tablet orally every 4 hours as needed for moderate-to-severe pain
- oxyCODONE 5 mg tab (OXYIR)
 - 1-2 tablet orally every 4 hours as needed for moderate-to-severe pain

Injectable

- morphine
 - 2-4 milligram intravenous push every 2 hours as needed for moderate-to-severe pain
- HYDROMorphine (DILAUDID)
 - 0.2 to 0.4 milligram intravenous push every 2 hours as needed for moderate-to-severe pain
- ketorolac (TORADOL)
 - 15 milligram intravenous push every 6 hours as needed for severe pain Contraindicated for use longer than 5 days

Patient-Controlled Analgesia

- For Patient controlled analgesia select the patient controlled analgesia order set

GI Side Effect Medications

Antiemetics

- promethazine (PHENERGAN)
 - 12.5-25 milligram orally every 6 hours as needed for nausea/vomiting potentially inappropriate in patients 65 years or older
 - 25 milligram rectally every 6 hours as needed for nausea/vomiting potentially inappropriate in patients 65 years or older
- metoclopramide (REGLAN)
 - 10 milligram orally every 6 hours as needed for nausea/vomiting
 - 10 milligram intravenously every 6 hours as needed for nausea/vomiting
- ondansetron (ZOFRAN)
 - 4 milligram intravenously every 4 hours as needed for nausea/vomiting

Stool Softeners/Laxatives (Recommended for use with Narcotics, EXCEPT if patient had bowel surgery)

- docusate sodium (COLACE)
 - 100 milligram orally 2 times a day - (HOLD for loose stools)
- bisacodyl (DULCOLAX)
 - 5 milligram orally 2 times a day - (HOLD for loose stools)
 - 10 milligram rectally once a day as needed for constipation - (HOLD for loose stools)

Initials _____

(place patient label here)

Patient Name: _____



Order Set Directions:

- > (√)- Check orders to activate; Orders with pre-checked box will be followed unless lined out.
- > Initial each place in the pre-printed order set where changes such as additions, deletions or line outs have been made
- > Initial each page and Sign/Date/Time last page

PROVIDER ORDERS

-
- polyethylene glycol 3350 17 gram oral powder packet (MIRALAX)
 - 17 milligram orally once a day IN 8 OZ (240 milliliters) OF WATER (HOLD for loose Stools)
 - sennosides-docusate sodium 8.6 mg-50 mg tab (SENOKOT-S)
 - 2 tablet orally once a day (HOLD for loose Stools)
 - magnesium hydroxide (MILK OF MAGNESIA)
 - 30 milliliter orally once a day, at bedtime as needed for constipation

Consults

- Pain Service Consult: Initiate pain management protocol

Provider Signature: _____ Date: _____ Time: _____