Patier Name		Benefis HEALTH SYSTEM Benefis Hospitals
Order Set D > > > >	rections: (√)- Check orders to activate; Orders with pre-checked box ☑ will be followed unless lined out. Initial each place in the pre-printed order set where changes such as additions, deletions or line outs have been made Initial each page and Sign/Date/Time last page	PROVIDER ORDERS
Diagnosis:		
ADUL [*]	ROUTINE ERCP ORDERS	Version 4 05/14/2014
 □ Consent patient for "moderate sedation, fluoroscopy, Endoscopic Retrograde CholangioPancreatogram, possible sphincterotomy, possible common bile duct stone removal, possible biopsy, possible biliary/pancreatic stent "with: □ JT Molloy MD or □ GD Spencer MD 		
	Contact GI lab (5224) for day and time for procedure.	
	☐ NPO (including dobhoff / NG tube feeds off) at least 6 hours prior to procedure.	
	Saline lock, if no IV currently running.	
	Levaquin 500 mg IV 30 minutes prior to procedure for known or suspected obstruction of biliary tree or synthetic vascular graft less than 1 year old. (see alternate below for Levaquin or quinolone allergy).	
	Merrem 1 gram IV 30 minutes prior to procedure for known or suspected obstruction of biliary tree <u>or</u> synthetic vascular graft less than 1 year old. (Use for patient allergic to quinolones; See alternate below for history of anaphylactic reaction to PCN or hypotension with PCN).	
	Clindamycin 600 mg IV <u>AND</u> Aztreonam 2 grams IV 30 minutes prior to procedure for known or suspected obstruction of biliary tree <u>or</u> synthetic vascular graft less than 1 year old. (Use for patient allergic to Levaquin <u>and</u> Merem or history of anaphylactic reaction to PCN or hypotension with PCN).	
	PT/INR, PTT if history of bleeding and no PT, PTT result done within last 7 days found on chart.	
	CBC if none done within last 7 days on chart.	

Other _____