	(place patient label here)	
Patie	ent Name:	
Order Set	= :: : : : : : : : : : : : : : : :	
>	(√)- Check orders to activate; Orders with pre-checked box ☑ will be foll	
>	Initial each place in the pre-printed order set where changes such as add Initial each page and Sign/Date/Time last page	ditions, deletions or li
 Diagnosi:	S:	



 I nitial each place in the pre-printed order set where changes such as additions, deletions or line outs have been made I nitial each page and Sign/Date/Time last page 	PROVI DER ORDERS
Diagnosis:	
Allergies with reaction type:	
Contrast Allergy Prophylaxis	Version 1 11/10/14
Hypersensitivity Prophylaxis Agents: For patients with a history of a dye SELECT ONE PREPROCEDURE OPTION Beginning 1 hour prior to procedure start time: Select all methylPREDNISolone 125 mg solution for injection (SOLUMEDROL) 125 milligram intravenously once one hour prior to procedure diphenhydrAMINE (BENADRYL) 50 milligram intravenously once 1 hour prior to procedure	allergy to iodine, contrast or
Beginning 13 hours prior to procedure start time: Select all predniSONE 50 milligram orally 13, 7 and 1 hour prior to procedure diphenhydrAMINE (BENADRYL) 50 milligram orally once 1 hour prior to procedure	
Beginning 3 days prior to procedure start time: Select all predniSONE 10 milligram orally 2 times a day x 2 days. First dose 2 days prior to day predniSONE	ay of procedure
□ 50 milligram orally 1 hour prior to procedurediphenhydrAMINE (BENADRYL)□ 50 milligram orally once 1 hour prior to procedure	