

Allergies with reaction type:

## **ICU Hypertonic Saline for Increased ICP Weaning Protocol**

## General

• After provider order for initiation of a protocol, nursing may place orders found within the protocol using the Policy/Protocol - No Esign Req order source.

Version 1 4/10/18

• Recommended for patient Age > 18 years old

## **Nursing Orders**

- ☑ **STOP** ICP Adjustment protocol
- ☑ When serum Na+ and serum osmolality results, notify provider immediately
- ☑ After 12 hours of *starting* ICP Weaning Protocol, draw Na+ and serum osmolality
- ☑ After 12 hours of *discontinuing* ICP Weaning Protocol, draw Na+ and serum osmolality

## **IV Fluids - Maintenance Specific Fluid**

<u>3% NaCl Infusion</u>	<u>Treatment</u>
60 mL/hr or GREATER	45 mL/hr for 8 hours, then 20 mL/hr for 8 hours, then 10 mL/hr for 8 hours, then stop
45 mL/hr-59 mL/hr	25 mL/hr for 8 hours, then 15 mL/hr for 8 hours, then 5 mL/hr for 8 hours, then stop
30 mL/hr-44 mL/hr	20 mL/hr for 8 hours, then 10 mL/hr for 8 hours, then 5 mL/hr for 8 hours, then stop
BELOW 30 mL/hr	10 mL/hr for 8 hours, then 5 mL/hr for 8 hours, then stop