

(place patient label here)

Patient Name: _____



PROVIDER ORDERS

Order Set Directions:

- > (✓) - Check orders to activate: Orders with pre-checked box will be followed unless lined out.
- > Initial each place in the pre-printed order set where changes such as additions, deletions or line outs have been made
- > Initial each page and Sign/Date/Time last page

Diagnosis: _____

Allergies with reaction type: _____

EDPED Rapid Sequence Intubation

Version 1 8/18/15

- This pediatric order set is intended for use in patients 1 month through 17 years of age and less than 50 kilograms

Nursing Orders

- Vital signs per unit standard
- Vital signs non unit standard _____
- Measure (NOT STATED) Weight in KG
- Cardiac monitor
- Pulse oximetry, continuous
- Continuous End Tidal CO2 Monitor

IV/ Line Insert and/or Maintain

- Peripheral IV insert/maintain

Medications

Pretreatment for RSI to prevent Bradycardia

atropine 1mg/10 mL syringe

- 0.02 milligram/kilogram intravenously once (Max 0.5 mg)
- 0.02 milligram/kilogram intramuscularly once (Max 0.5 mg)

- Optional for increased ICP

lidocaine (PF) 100 mg/5 mL (2 %) injection syringe

- 1.5 milligram/kilogram intravenously once (Max 100 mg)

Analgesics

fentaNYL 100 mcg/2 mL ampule

- ____ microgram intravenously as needed for pain during procedure; may repeat every 1-3 minutes to desired effect

- (Recommended dose range 0.5-1.5 microgram/kilogram)

Anesthetics / Sedatives

- Avoid the use of etomidate for rapid sequence intubation in patients with severe sepsis

etomidate (AMIDATE) 40 mg/20 mL vial

- 0.3 milligram/kilogram over 30-60 seconds intravenously once

- Use with caution for patients with increased ICP

ketamine (KETALAR)

- ____ milligram over 1-2 minutes intravenously once

(Recommended dose range 1-2 milligram/kilogram)

midazolam (VERSED) 5 mg/5mL vial

- ____ milligram intravenously once (Max 2 mg)

(Recommended dose range 0.02-0.03 milligram/kilogram)

Neuromuscular Blocking Agents

succinylcholine 200mg/10 mL vial

- 1 milligram/kilogram intravenously once

rocuronium (ZEMURON)

- 1 milligram/kilogram (using ideal body weight) intravenously once

vecuronium (NORCURON) 10 mg vial

- 0.15 milligram/kilogram intravenously once

Laboratory

- Blood gas study [] Arterial [] Venous [] Capillary

Radiology

- XR Chest Single, portable, Reason for exam: Verify ETT placement

Provider Signature: _____ Date: _____ Time: _____