

(place patient label here)

Patient Name: _____



PROVIDER ORDERS

Order Set Directions:

- > (✓)- Check orders to activate; Orders with pre-checked box will be followed unless lined out.
- > Initial each place in the pre-printed order set where changes such as additions, deletions or line outs have been made
- > Initial each page and Sign/Date/Time last page

Diagnosis: _____

Allergies with reaction type: _____

ICU Hypertonic Saline for Increased ICP Weaning Protocol

Version 1 4/10/18

General

- After provider order for initiation of a protocol, nursing may place orders found within the protocol using the Policy/Protocol - No Esign Req order source.
- Recommended for patient Age > 18 years old

Nursing Orders

- STOP** ICP Adjustment protocol
- When serum Na+ and serum osmolality results, notify provider immediately
- After 12 hours of *starting* ICP Weaning Protocol, draw Na+ and serum osmolality
- After 12 hours of *discontinuing* ICP Weaning Protocol, draw Na+ and serum osmolality

IV Fluids - Maintenance Specific Fluid

<u>3% NaCl Infusion</u>	<u>Treatment</u>
60 mL/hr or GREATER	45 mL/hr for 8 hours, then 20 mL/hr for 8 hours, then 10 mL/hr for 8 hours, then stop
45 mL/hr-59 mL/hr	25 mL/hr for 8 hours, then 15 mL/hr for 8 hours, then 5 mL/hr for 8 hours, then stop
30 mL/hr-44 mL/hr	20 mL/hr for 8 hours, then 10 mL/hr for 8 hours, then 5 mL/hr for 8 hours, then stop
BELOW 30 mL/hr	10 mL/hr for 8 hours, then 5 mL/hr for 8 hours, then stop