(place patient label here) Patient Name: Order Set Directions: > (√)- Check orders to activate; Orders with pre-checked box ☑ will be followed unless lined out. > Initial each place in the pre-printed order set where changes such as additions, deletions or line outs have been made > Initial each page and Sign/Date/Time last page	Benefis HEALTH SYSTEM Benefis HOSPITALS PROVI DER ORDERS
Diagnosis:	
Allergies with reaction type:	
ED Common IV Drips	Version 1 8/18/15
Sedative Infusions propofol (DIPRIVAN)10 mg/mL 5 microgram/kilogram per minute continuous intravenous infusion Mamicrogram/kilogram per minute to achieve appropriate sedation leve dexmedetomidine (PRECEDEX) in NS 400 mcg/100 ml (4 mcg/ml) 0.5 microgram/kilogram per hour continuous intravenous infusion; Mamicrogram/kilogram per hour to achieve appropriate sedation level	S) DC prior to extubation
Vasoactive Infusions diltiazem (CARDIZEM) 125 mg in 125 mL □ 5 milligram/hour continuous intravenous infusion titrate to keep HR < DOBUTamine (DOBUTREX) [500 milligrams/ 250 milliliters D5W] □ 2.5 microgram/kilogram per minute continuous intravenous infusion with a mean arterial pressure greater than or equal to 65 mmHg and (MAX 10 microgram/kilogram per minute) DOPamine in D5W 400 mg/250 mL (1,600 mcg/mL) IV □ 5 microgram/kilogram per minute continuous intravenous infusion: ti	: titrate to keep SBP > 90 mmHg I Heart Rate < 140 beats per minute Itrate to keep SPB greater than or
equal to 90mmHg and MAP greater than or equal to 65 mmHg (MAX EPINEPHrine in Dextrose 5% water (0.016 milligram/milliliter) 0.5 microgram/kilogram per minute continuous intravenous infusion microgram/kilogram per minute) Titrate to keep systolic blood pressuniCARdipine (CARDENE) 20 mg in 200 mL 5 milligram/hour continuous intravenous infusion titrate to keep SBP	(dose range 0.1-1 ure greater than 90 mmHg
or equal to 65 mmHg (MAX 15 milligram/hour) norepinephrine bitartrate in normal saline 4 mg/250 mL IV (LEVOPHED) □ 1 microgram/minute continuous intravenous infusion: titrate to keep greater than or equal to 65 mmHg (MAX 30 microgram/minute) phenylephrine in NS (preserv free) 20 mg/250 mL (80 mcg/mL) IV (NEO-S □ 20 microgram/minute continuous intravenous infusion titrate to keep	YNEPHRINE)

MAP greater than or equal to 65 mmHg (MAX 180 microgram/minute)

unit/minute)

vasopressin (PITRESSIN)[40 unit/100 milliliter NS]

□ 0.03 unit/minute continuous intravenous infusion titrate to keep MAP greater than 80 mmHg (MAX 0.04