


Neonatal Sepsis Screening

(Place Patient Label Here)

NURSING



- SIGN OF NEONATAL SEPSIS**
- Respiratory Distress: GFR, apnea, tachypnea (Resting RR > 60), requiring O2 support
 - Temperature instability: Temp > 99.4 or < 97.2
 - Feeding intolerance: Vomiting, abdominal distention, poor feeding pattern
 - Abnormal skin perfusion: Mottling, pale or gray color, capillary refill > 3 seconds
 - Abnormal heart rate or BP: Tachycardia (Resting HR > 165), bradycardia (Resting HR < 100), hypotension (Resting SBP < 55 or MBP < 35)
 - Abnormal neurologic status: Lethargy, hypotonia, irritability, seizures

- INTRAPARTUM GBS PRPHYLAXIS INDICATORS**
- Previous infant with invasive B=GBS disease
 - GBS bacteriuria during any trimester of current pregnancy
 - Positive GBS vaginal-rectal screening in late gestation (33-37 weeks) during current pregnancy
 - Unknown GBS status at the onset of labor (culture not done, incomplete or results unknown) and any of the following
 - *Delivery at < 37 weeks gestation
 - *Amniotic membrane rupture ≥ 18 hours
 - *Intrapartum temperature ≥ 100.4F

PROVIDER

Provider transfers infant to NICU For full diagnostic evaluation and antibiotic therapy

Provider orders blood culture and CBCD and evaluate lab results and order additional labs or transfer to NICU if full diagnostic workup is indicated