

(place patient label here)

Patient Name: _____



PROVIDER ORDERS

Order Set Directions:

- > (✓) - Check orders to activate: Orders with pre-checked box will be followed unless lined out.
- > Initial each place in the pre-printed order set where changes such as additions, deletions or line outs have been made
- > Initial each page and Sign/Date/Time last page

Diagnosis: _____

Allergies with reaction type: _____

Wound Care Protocol

Version 4 11/10/14

- After provider order for initiation of wound care protocol, nursing may place the following orders using the per protocol order source.

General

- Apply Elta Moisturizer to Dry Skin as needed
- Rooke Boots as needed for Lower Extremity Ulcer Treatment or Prevention
- Specialty Air Mattress Bed as needed for Patients with Stage 3-4 Pressure Ulcers
- Chair Cushion

Skin Tear/ Open Blister/ Abrasion Care

- Location: _____
- Cleanse and Irrigate Area with Saline
- Apply No Sting /Prep to Surrounding Intact Skin
- Approximate Edges if able and Apply Steri-strips. Change prn.
- Apply Tubegrip to Arms as needed for Compression

Apply Dressing Types as Indicated

For Fragile Skin and/or Draining Wounds:

- Oil emulsion and wrap with gauze- Change QD.

For Non-fragile Skin:

- Foam and Silicone size < 6cm- Change Q3D
- Foam and Tegaderm size > 6cm- Change Q3D

Incontinence/ Moisture Associated Skin Damage

- Location: _____
 - Cleanse with Ready Bath
 - Apply Thin Layer of Calazime to Area QID and prn
 - May Use Peripad with Mesh Pants- No Briefs
 - Place Stockinett to Skin Folds for Absorption prn
- miconazole powder (LOTRIMIN)
- 1 application topically 2 times a day as needed for bright red or purple rash with or without satellite lesions

Apply Dressings Types as Indicated

For Skin Loss with Minimal Drainage:

- Foam and Silicone size < 6cm- Change Q3D
- Foam and Tegaderm size > 6cm- Change Q3D

For Skin Loss with Moderate Drainage:

- Aquacel and Foam and Wrap/Tape to Secure- Change QD

Pressure Ulcer/ Extremity Ulcer Care

- Location: _____
- Cleanse and Irrigate with Saline
- Apply No Sting /Prep to Surrounding Intact Skin

Apply Dressings Types as Indicated

Dry Crusted Ulcers on Lower Extremity ONLY

- Betadine Swab BID

Ulcers with Pink or Red Ulcer Base

For Partial Thickness Ulcers with Minimal Depth:

- Foam and Silicone size < 6cm- Change Q3D
- Foam and Tegaderm size > 6cm- Change Q3D

(place patient label here)

Patient Name: _____



PROVIDER ORDERS

Order Set Directions:

- > (✓)- Check orders to activate: Orders with pre-checked box will be followed unless lined out.
- > Initial each place in the pre-printed order set where changes such as additions, deletions or line outs have been made
- > Initial each page and Sign/Date/Time last page

- Thin Duoderm where flexibility is needed- Change Q3D

For Partial Thickness Ulcers with Depth:

- Aquacel (moisten if dry) and Foam and Silicone size < 6cm- Change QOD
- Aquacel (moisten if dry) and Foam and Tegaderm size > 6cm- Change QOD

For Partial Thickness Dry Ulcers:

- Hydrogel and Foam and Silicone size < 6cm- Change QOD
- Hydrogel and Foam and Tegaderm size > 6cm- Change QOD

For Partial Thickness Sacral Ulcers:

- Sacral Foam and Silicone- Change QOD

For Full Thickness Dry or Moist Ulcers:

- Aquacel (moisten if dry) and Foam and Silicone size < 6cm- Change QOD
- Aquacel (moisten if dry) and Foam and Tegaderm size > 6cm- Change QOD

Ulcers with Necrotic Tissue or Surrounding Erythema or Edema

For Partial or Full Thickness Ulcers With or Without Drainage:

- Aquacel AG (moisten if dry) and Foam and Wrap/Tape to Secure- Change QD

For Non-tunneling Ulcers With Dry Necrotic Tissue:

- Hydrogel and Foam and Wrap/Tape to Secure- Change QD

For Full Thickness Ulcers With Tunneling:

- Saline Moistened Fluffed Gauze and Gauze and Wrap/Tape to Secure- Change Q8H.

For Partial Thickness Dry Ulcers SELECT:

cadexomer iodine (IODOSORB 0.9% GEL)

- 1 application topically once a day and Foam and Wrap/Tape to Secure