

(place patient label here)

Patient Name: _____



PROVIDER ORDERS

Order Set Directions:

- > (✓)- Check orders to activate; Orders with pre-checked box will be followed unless lined out.
- > Initial each place in the pre-printed order set where changes such as additions, deletions or line outs have been made
- > Initial each page and Sign/Date/Time last page

Diagnosis: _____

Allergies with reaction type: _____

Bariatric Surgery Post Op Day 1

Version 2 Approved 11/13/2017

Nursing Orders

- BVI
- Perform Methylene Blue Test, if ordered
- Discontinue JP Drain
- Wound Care and Dressing Instructions for Bariatric Surgery Post Op
- Patient Education: Teach patient drain care and to log output over each 24 hours
- Notify Physician when discharge criteria met for Bariatric Surgery patients:
 - Tolerating 4-6oz PO fluids per hour, which includes the protein shake fluid intake
 - Tolerating protein supplement at a rate consistent with at least 50-60 grams of protein per day
 - Nausea controlled adequately with oral antiemetics to allow adequate PO fluid tolerance
 - Pain control with oral analgesics
 - Ambulating adequately to full ADLs
 - Urinating adequately

Diet (pick only one)

- Clear Liquid Diet with Bariatric Modifiers
- Full Liquid Diet with Bariatric Modifiers
- Optisource/protein supplement (with Full Liquid Diet) 4 ounces every 2 hours, 6 times per day
Additional Instructions: No red dye, no straws, no caffeine, no carbonated beverage, no juices. Artificial sweeteners ONLY.
Required intake: 6 ounces fluid per hour.

IV/ Line Insert and/or Maintain

- Convert Peripheral IV to Saline Lock when tolerating PO

Medications

- pantoprazole IV
 - 40 milligram intravenously once
- pantoprazole Oral
 - 40 mg orally every morning
- methylene Blue
 - 3 milliliters to 30ml of ice chips

Analgesics: Oral

- acetaminophen 500 mg tablet (TYLENOL)
 - 1,000 milligram orally every 6 hours
- oxyCODONE 5 mg tablet
 - 1-2 tablets orally every 4 hours as needed for moderate-to-severe pain
- HYDROcodone/acetaminophen 5/325mg tablet (NORCO)
 - 1-2 tablets orally every 4 hours as needed for moderate pain

Provider Signature: _____ Date: _____ Time: _____